

My Colonoscopy Prep Calendar—Bowel Prep using SUPREP



7 days
before
procedure

Your Pharmacy Shopping List:

- One bottle Milk of Magnesia (or generic equivalent of magnesium hydroxide).
- Prescription for the SUPREP.



7 days
before
procedure

Your Grocery Shopping List:

- Purchase clear liquid products—examples include:
 - sports drinks (not red, blue or purple)
 - apple or white grape juice
 - ginger ale
 - clear broth or bouillon
 - water
 - coffee/tea without milk
 - clear carbonated beverages
 - plain Jell-O (not red, blue or purple and no fruit or toppings)
 - Italian ice



3 days
before
procedure

- STOP eating all fruits, vegetables, seeds and nuts.
- Stop taking any iron or fiber supplements.
- Confirm ride and contact phone number for your colonoscopy. You **MUST** have a ride home from a responsible adult as you will have received sedation for the procedure. (**NO** taxis, Lyft, Uber, etc).
- Please inform the person providing your ride that the appointment will take approximately 2 ½-3 hours.
- If you are **unable** to make your appointment, please call: **603.665.2470**.



2 days
before
procedure

- Only consume **CLEAR LIQUIDS THE ENTIRE DAY** (see 7 days list for suggested liquids).
- DO NOT** eat any solid food.
- It is important that you drink lots of fluids 2 days before your procedure as the preparation is very dehydrating.
- Take 2 tablespoons of Milk of Magnesia at dinner time.



1 day
before
procedure

- Consume only **CLEAR LIQUIDS** the entire day (no solid food, dairy, juice with pulp, alcohol or anything red, blue or purple in color).
- In the morning:** Prepare the solution of SUPREP and **refrigerate**.
- 4 PM:**
- Drink one of the 6 oz. bottles of SUPREP and **refrigerate**.



procedure day

- Have **NOTHING to eat or drink after midnight**, except for finishing the SUPREP and/or taking any medications as ordered.
- 5 hours before your check-in time:** FINISH the SUPREP by drinking the second 6 oz. bottle.

Important:

- If you are on **blood pressure medication**, you must **take your medication the morning** of your procedure, prior to check-in.

Thank you for your consideration.



My Colonoscopy Prep Calendar—Bowel Prep using SUPREP

IMPORTANT information

Please continue your daily medication regimen except for:

- Diabetic Patients:** Do not take any diabetes medication on the day of your procedure. If you have any questions, please call your prescribing provider. If you take insulin, call your prescribing provider and ask for instructions.
- Patients on Blood Thinners: YOU MUST** check in with your prescribing physician before stopping your medication. If you are currently taking an ASPIRIN daily, do not stop. Below is a list of common blood thinners and when, with **physician approval**, to discontinue them:
 - Ticlid (ticlopidine)—stop 10 days prior to procedure
 - Effient (prasugrel)—stop 7 days prior to procedure
 - Coumadin (warfarin), Plavix (clopidogrel) or Brilinta—stop 5 days prior
 - Pradaxa (dabigatran), Eliquis (apixaban) or Xarelto (rivaroxaban)—stop 48 hours prior
- Patients on Erectile Dysfunction medication:** Viagra, Cialis, Adcirca, Sildenafil or Revatio—stop 24 hours prior.
- Patients on Appetite Suppressant medication:** phentermine (brand names Lomaira, Adipex-P; also as phentermine/topiramate in Qsymia)—stop 4 days prior to procedure.
- Please **bring a list of your current medication** to the procedure and include dosing information.
- You will be contacted by Endoscopy **one week prior** to your procedure to review important details. You will also be contacted 48 hours prior to the procedure to be given your arrival time.
- If you have any discomfort** around the perianal region, use wet wipes instead of toilet paper and apply Vaseline or A&D Ointment to the rectum.
- If you **are unable to make your appointment**, please call: **603.665.2470**. Thank you for your consideration.

Diabetic and Weight Loss Medications

- ❑ **If you are taking the following medications, you must ensure that your procedure takes place at least 8 days after your most recent dose:**

Generic Drug	Brand Name
Dulaglutide	Trulicity
Exenatide (ER)	Bydureon BCise
Semaglutide	Ozempic, Wegovy, others
Tirzepatide	Mounjaro, Zepbound
Albiglutide	Tanzeum, Eperzan

- ❑ **If you are taking the following medication, it will need to be held for 7 days prior to your procedure:**

Generic Drug	Brand Name
Exenatide (IR)	Byetta
Liraglutide (3mg)	Saxenda
Liraglutide (1.2mg, 1.8mg)	Victoza
Lixisenatide	Adlyxin
Semaglutide	Rybelsus

- ❑ **Please ensure you are checking your blood sugar levels at a more frequent rate while you are holding these medications.**